

CLINICAL ASSESSMENT OF TUBERCUL (IGRA)

H TB IGRA (I)

3. I G R A (IGRA)

____ / ____ / ____ (____), E____, ____
L / /

____ (____)

____ / ____ / ____ (____), E____, ____
L / /

____ (____)

4. C : (R | TST IGRA)

____ / ____ / ____
L / /

M | P TST IGRA

All students with a positive TST or IGRA with no signs of active disease on chest x-ray should receive a recommendation to be treated with medication. However, students in the following groups are at increased risk of progression from LTBI to TB disease and should be treated as possible.

- Infected with HIV
- Recently infected with M. tuba (y inJ_plAllatment a)0.9 (8K(wit in the p)0.5 a)0.5 (s)-4.9 (2 y)71.2 (a)0.5 (r)142.3 sy)
- Hiestoreated(ormint)0.6 dequactlreateo TB dasetudtest-rai(og)1.9 (r)-6 (a)0.5 ptent with prio TB dasease
- Recivwingimmunosup pressive thprpv u stumioonector61.7 -alpha_ (NFd)antfss(, ya)-5.4 (s)-4.9 (t)0.8 (micn (or)--1.1 ticoes)-4. (pr)-2.9 edni(oneg p)0.5 e(d(a)-5.4 ()4)0.5 ,d(ormmmunosupp)0.5 (pr)31 (essiv)5.5 (ed(r)12.9 uge th(r)-6 (pv)5.4 (y f)-1.1 (ollo)6
- Diagnos(ed withsilor61.n9 0d)065 (aab)0.5 (et)0.8 (rs mellitu9 0c)6.1 (hr)-2.9 (oncn r)-3.1 elnalfrmicad(ormca)0.5 (nb)0.5 ((dof t
- _____ S71.7 (tud)0.6 (ent ag)1.9 (r)-3 (ers t)1.8 (o r)-3 (eceiv)5.5 (e tr)-3.1 elatment
- _____ S71.7 (tud)0.6 (ent declines tr)-3 (ea)0.5 (tment at t is time*)]TJ 0 Tc 0 -5.9 Td (H)Tj -0.001 Tc 0.467 0 Td [(ea)0.6 (lthcar)-
- _____

Healthcare Prvda(dSignatur)-2.9 ed: _____ Datr: _____

Prtvda(dA)-4.6 dedress: _____



FORM MUSAMPED BY HEALTH CARE OFFICE OR IT WILL NOCCEPTED.

*Pecommatioes setforthsb by the American College HealthdAssociatioe, ano declines treatment will be asko repeat a0chay every two years while enry University